

GUIDELINE 2b Skin Infections in Wrestling

July 1981 • Revised June 2003 _

Data from the NCAA Injury Surveillance System (ISS) indicate that skin infections are associated with at least 15 percent of the practice time-loss injuries in wrestling. It is recommended that qualified personnel examine the skin over the entire body, and the hair of the scalp and pubic areas of all wrestlers before any participation in the sport.

Open wounds and infectious skin conditions that cannot be adequately protected to prevent their exposure to others should be considered cause for medical disqualification from practice or competition.

Categories of skin conditions and examples include:

- 1. Bacterial skin infections
 - a. impetigo;
 - b. erysipelas;
 - c. carbuncle;
 - d. staphylococcal disease;
 - e. folliculitis (generalized);
 - hidradentitis suppurativa;
- 2. Parasitic skin infections
 - a. pediculosis;
 - b. scabies:
- 3. Viral skin infections
 - a. herpes simplex;
 - b. herpes zoster (chicken pox);
 - c. molluscum contagiosum; and

4. Fungal skin infections—tinea corporis (ringworm).

Note: Current knowledge indicates that many fungal infections are easily transmitted by skin-to-skin contact. In most cases, these skin conditions can be covered with a securely attached bandage or nonpermeable patch to allow participation.

Besides identification of infected individuals and their prompt treatment, prevention can be aided through proper routine cleaning of all equipment, including mats and shared common areas, such as locker rooms.

This guideline is intended for general information only. Team physicians, athletic trainers, coaches and others who work directly with wrestling should refer to the current year's NCAA Wrestling Rules Book or Wrestling Championships Handbook for rules regarding specific skin infections.

References

- 1. Belongia EA, Goodman JL, Holland EJ, et. al.: An outbreak of herpes gladiatorium at a high school wrestling camp. <u>The New England Journal of Medicine</u>. 325(13):906-910, 1991.
- 2. Mast, E. and Goodman, R.: Prevention of Infectious Disease Transmission in Sports. SportsMedicine 24(1):1-7,1997.
- 3. Dorman, JM.: Contagious diseases in competitive sport: what are the risks?

 <u>Journal of American College Health</u>
 49(3):105-9, 2000 Nov.
- **4.** Adams, BB.: Transmission of cutaneous infection in athletics. British Journal of Sports Medicine 34(6):413-4, 2000 Dec.
- 5. Beck, CK.: Infectious diseases in sports: Medicine and Science in Sports and Exercise 32(7 Suppl):S431-8, 2000 Jul.
- **6.** Lindenmayer JM, Schoenfeld S, O'Grady R, Carney JK.: Methicillin-resistant

- Staphylococcus aureus in a high school wrestling team and the surrounding community. <u>Archives of Internal Medicine</u> 158(8):895-9, 1998 Apr.
- 7. Kohl TD, Martin DC, Nemeth R, Hill T, Evans D.: Fluconazole for the prevention and treatment of tinea gladiatorum.

 Pediatric Infectious Disease Journal
 19(8):717-22, 2000 Aug.
- 8. Anderson BJ.: The Effectiveness of Valacyclovir n Preventing Reactivation of Herpes Gladiatorum in Wrestlers. Clin J Sports Med 9(2):86-90, 1999 Apr.
- 9. Vasily DB, Foley JJ.: More on Tinea Corporis Gladiatorum. <u>J Am Acad</u> <u>Dermatol</u> 2002, Mar.
- **10.** Vasily DB, Foley JJ, First Episode Herpes Gladiatorum: Treatment with Valacyclovir. (manuscript submitted for publication).